

Betti numbers are testable ^{*}

Gábor Elek

July 30, 2009

Abstract

We prove that the Betti numbers of simplicial complexes of bounded vertex degrees are testable in constant time.

1 Introduction

Property testing in bounded degree graphs was introduced in the paper of Goldreich and Ron [4]. In this paper we study property testing for bounded degree simplicial complexes in higher dimensions. Let $d \geq 2$ be a natural number and consider finite simplicial complexes where each vertex (zero dimensional simplex) is contained in at most d edges (1-dimensional simplex). Of course, such a complex can be at most d -dimensional. What does it mean to test the p -th Betti number of such a simplicial complex? First fix a positive real number $\varepsilon > 0$. A tester takes a simplicial complex K as an input and pick $C(\varepsilon)$ random vertices. Then it looks at the $C(\varepsilon)$ -neighborhoods of the chosen vertices. Based on this information the tester gives us a guess $\hat{b}^p(K)$ for the p -th Betti number $b^p(K)$ of the simplicial complex such a way that :

$$\text{Prob}\left(\frac{|\hat{b}^p(K) - b^p(K)|}{|V(K)|} > \varepsilon\right) < \varepsilon,$$

where $V(K)$ is the set of vertices in K . In other words, we can estimate the p -th Betti number very effectively with high probability knowing only a small (random) part of the simplicial complex. The goal of this paper is to show the existence of such a tester for any $\varepsilon > 0$. That is to prove the following theorem:

Theorem 1 *Betti-numbers are testable for bounded degree simplicial complexes.*

For graphs the 0-th Betti number is just the number of components and the first Betti number can be computed via the 0-th Betti number and the Euler-characteristic, hence it is not hard to see that such tester exists. For connected surfaces one can also calculate the first Betti number using just the number of

^{*}Research sponsored by OTKA Grant K 69062 and 49841

vertices, edges and triangles. However in higher dimensions there is no such formula even for triangulated manifolds. Note that this paper was not solely motivated by the paper of Goldreich and Ron, but also by the solution of the Kazhdan-Gromov Conjecture by Wolfgang Lück [5]. The workhorse lemma of our paper is basically extracted from his paper using a slightly different language. It is very important to note that our proof works only for Betti numbers of real coefficients and we do not claim anything for the Betti numbers of mod- p coefficients.

2 The convergence of simplicial complexes

Let Σ^d be the set of finite simplicial complexes K of vertex degree bound d that is any 0-dimensional simplex is contained in at most d 1-dimensional simplices. We denote by K_i the set of i -simplices in K and by G_K the 1-skeleton of K , that is $V(G_K) = K_0$, $E(G_K) = K_1$. A rooted r -ball of degree bound d is a simplex $L \in \Sigma^d$ with a distinguished vertex x such that for any $y \in V(G_L)$, $d(x, y) \leq r$, where $d(x, y)$ is the shortest path distance of x and y in the graph G_L . We denote by $Z^{r,d}$ the rooted isomorphism classes of rooted r -balls. If $K \in \Sigma^d$ and $p \in V(K)$ then let $G_r(p)$ be the rooted r -ball in the 1-skeleton G_K and $B_r(p)$ is the set of simplices σ such that all vertices of σ are in $G_r(p)$. Then $B_r(p)$ is a rooted r -ball of vertex degree bound d .

For $\alpha \in Z^{r,d}$ we denote by $T(K, \alpha)$ the set of vertices p such that $B_r(p) \cong \alpha$. We set

$$p_K(\alpha) := \frac{|T(K, \alpha)|}{|K_0|}.$$

We say that $\{K^n\}_{n=1}^\infty \subset \Sigma^d$ is convergent (see [1] for the graph case) if

- $|K_0^n| \rightarrow \infty$
- $\lim_{n \rightarrow \infty} p_{K_n}(\alpha)$ exists for any $r \geq 1$ and $\alpha \in Z^{r,d}$.

It is easy to see that any sequence $\{L^n\}_{n=1}^\infty \subset \Sigma^d$ such that $|L_0^n| \rightarrow \infty$ contains a convergent subsequence.

Let $\alpha_1, \alpha_2, \alpha_3, \dots$ be an enumeration of all the r -balls, $r \geq 1$. Then we have the pseudo-metric

$$d_s(K, L) := \sum_{i=1}^{\infty} \frac{1}{2^i} |p_K(\alpha_i) - p_L(\alpha_i)|.$$

Clearly, if $\{K^n\}_{n=1}^\infty$ be an increasing sequence of simplicial complexes they are Cauchy if and only if they are convergent. By an oriented d -complex Q we mean an element of Σ^d equipped with a fixed orientation for each of its simplex. Note that we do not assume that the orientations are compatible in any sense. We denote by $\hat{\Sigma}^d$ the set of all finite oriented d -complexes. We also define oriented r -balls, the set $\hat{Z}^{r,d}$ of all oriented r -ball isomorphism classes and the probabilities $p_Q(\beta)$ accordingly. Naturally we can define the convergence of oriented d -complexes as well.

Proposition 2.1 *Let $\{K^n\}_{n=1}^\infty \subset \Sigma^d$ be a convergent sequence of d -complexes. Then one has an oriented copy Q^n for each K^n such that $\{Q^n\}_{n=1}^\infty \subset \hat{\Sigma}^d$ is convergent as well.*

Proof. Consider i.i.d random variables $\theta(x)$ distributed uniformly on $[0, 1]$. Let $(a_0, a_1, \dots, a_i) \in K_i^n$ be a positive orientation if

$$\theta(a_0) < \theta(a_1) < \dots < \theta(a_i).$$

Then the resulting sequence $\{Q^n\}_{n=1}^\infty \subset \hat{\Sigma}^d$ is convergent with probability 1. For details of the simple argument see ([3], Proposition 2.2) ■

Finally, we need a technical definition that we use in the subsequent sections. Let us consider the set Q_i of i -simplices in an oriented d -complex Q . If $\sigma, \tau \in Q_i$ we say that σ and τ are adjacent if they have at least one joint vertex. This way we can define the shortest path distance $d_i(\sigma, \tau)$ (where $d_i(\sigma, \tau) = \infty$ is possible). The ball $B_r^i(\sigma)$ is the set of i -simplices τ such that $d_i(\sigma, \tau) \leq r$. As above, we can define the classes $\hat{Z}_i^{r,d}$ and the sampling probabilities $p_{Q_i}(\beta)$. Note that if $\alpha, \beta \in \hat{Z}_i^{r,d}$ we say that α is isomorphic to β if they are isomorphic as simplicial complexes not only as metric spaces.

That is $p_{Q_i}(\beta)$ defined the following way. Let $\beta \in \hat{Z}_i^{r,d}$. Denote by $T(Q_i, \beta)$ the number of i -simplices τ such that the simplicial complex $B_r^i(\sigma)$ is isomorphic to β , where the isomorphism preserves the root-simplex. Then

$$p_{Q_i}(\beta) := \frac{|T(Q_i, \beta)|}{|Q_i|}.$$

Clearly, if $\{Q^n\}_{n=1}^\infty \subset \hat{\Sigma}^d$ is a convergent sequence then for any $\beta \in \hat{Z}_i^{r,d}$, $\lim_{n \rightarrow \infty} p_{Q_i^n}(\beta)$ exists.

3 Betti numbers and combinatorial Laplacians

Let $Q \in \hat{\Sigma}^d$ be an oriented simplicial complex. Let $C^i(Q)$ denote the euclidean space of real functions on the i -simplices of Q . Let us consider the cochain-complex

$$C^0(Q) \xrightarrow{d_0} C^1(Q) \xrightarrow{d_1} \dots$$

Recall that if $f \in C^q(Q)$ then

$$\begin{aligned} df(a_0, a_1, \dots, a_{q+1}) &= f(a_1, a_2, \dots, a_q) - f(a_0, a_2, \dots, a_{q+1}) + \dots \\ &\quad + (-1)^{q+1} f(a_0, a_1, \dots, a_q). \end{aligned}$$

Then $b^i(Q) = \dim \text{Ker } d_i - \dim \text{Im } d_{i-1}$ are the Betti numbers of Q . Note that they do not depend on the choice of the orientation of Q only the underlying simplicial complex.

The combinatorial Laplacians (see e.g. [2]) $\Delta_Q^i : C^i(Q) \rightarrow C^i(Q)$ are defined as

$$\Delta_Q^i := d_{i-1} d_{i-1}^* + d_i^* d_i.$$

The operators Δ_Q^i are positive and self-adjoint. Also,

$$\dim \text{Ker} \Delta_Q^i = b^i(Q).$$

Let us remark that by Lemma 2.5 of [2] we have the following information on the combinatorial Laplacians:

- $\Delta_Q^i(\sigma, \tau) \neq 0$ only if $\sigma = \tau$ or σ and τ are adjacent.
- $\Delta_Q^i(\sigma, \tau)$ is always an integer.
- $|\Delta_Q^i(\sigma, \tau)| \leq d + 1$.

4 Weak convergence of probability measures

First recall the notion of weak convergence of probability measures. Let $\{\mu_n\}_{n=1}^\infty$ be probability measures on the interval $[0, K]$. Then $\{\mu_n\}_{n=1}^\infty$ weakly converges to μ if for any continuous function $f \in C[0, K]$

$$\int_0^K f d\mu_n \rightarrow \int_0^K f d\mu.$$

For an example, let $\mu_n(\frac{1}{n}) = 1$, then the measures μ_n converge to the measure concentrated at the zero. Note that in this case $\lim_{n \rightarrow \infty} \mu_n(0) \neq \mu(0)$.

Now let

$$c(\mu_n) = \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^K \log \lambda d\mu_n.$$

The following theorem can be extracted from [5], nevertheless we provide a proof using only the language of real analysis, avoiding any reference to operators.

Theorem 2 *Suppose that $\{\mu_n\}_{n=1}^\infty$ weakly converges to μ and for any $n \geq 1$, $c(\mu_n) \geq 0$. Then $\lim_{n \rightarrow \infty} \mu_n(0) = \mu(0)$.*

Proof. First we need some notations. For a monotone function f ,

$$f^+(\lambda) = \inf_{\varepsilon \rightarrow 0} f(\lambda + \varepsilon)$$

For the measures μ_n let σ_n be their distribution function that is

$$\sigma_n(\lambda) = \mu_n([0, \lambda]).$$

Also, let $t(\lambda) = \mu([0, \lambda])$. Note that $\sigma_n^+ = \sigma_n$. Let

$$\overline{\sigma}(\lambda) := \limsup_{n \rightarrow \infty} \sigma_n(\lambda)$$

and

$$\underline{\sigma}(\lambda) := \liminf_{n \rightarrow \infty} \sigma_n(\lambda).$$

The following lemma trivially follows from the definitions.

Lemma 4.1 *Let f be a continuous function such that*

$$\chi_{[0, \lambda]}(x) \leq f(x) \leq \chi_{[0, \lambda + \frac{1}{k}]}(x) + \frac{1}{k}$$

for any $0 \leq x \leq K$, then

$$\sigma_n(\lambda) \leq \int_0^K f(\lambda) d\mu_n \leq \sigma_n(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

$$t(\lambda) \leq \int_0^K f(\lambda) d\mu \leq t(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

Proposition 4.1 $\bar{\sigma}(\lambda) \leq t(\lambda) = \underline{\sigma}^+(\lambda) = \bar{\sigma}^+(\lambda).$

Proof. By this lemma,

$$\bar{\sigma}(\lambda) \leq t(\lambda) \leq \bar{\sigma}(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

Hence $\bar{\sigma}(\lambda) \leq t(\lambda) = \underline{\sigma}^+(\lambda) \leq \bar{\sigma}^+(\lambda)$. Since $t(\lambda)$ is monotone, we have that $\bar{\sigma}(\lambda + \epsilon) \leq t(\lambda + \epsilon)$ and $\bar{\sigma}^+(\lambda) \leq t^+(\lambda) = t(\lambda)$. Thus our proposition follows. $\blacksquare$

The following elementary analysis lemma is proved in [6].

Lemma 4.2 *Let f be a continuously differentiable function on the positive reals and μ be a probability measure on the $[0, K]$ interval. Suppose that F is the distribution function of μ , that is $\mu[0, \lambda] = F(\lambda)$. Then for any $0 < \epsilon \leq K$:*

$$\int_\epsilon^K f(\lambda) d\mu = - \int_\epsilon^K f'(\lambda) F(\lambda) d\lambda + f(K)F(K) - f(\epsilon)F(\epsilon).$$

Assume that $K \geq 1$, then by the previous lemma

$$c(\mu_n) = \log K - \log(\epsilon) \sigma_n(\epsilon) - \int_\epsilon^K \sigma_n(\lambda) \frac{1}{\lambda} d\lambda.$$

Since

$$\int_\epsilon^K \frac{\sigma_n(0)}{\lambda} = (\log K - \log(\epsilon)) \sigma_n(0),$$

we have that

$$c(\mu_n) = \log K (1 - \sigma_n(0)) - \int_0^K \frac{\sigma_n(\lambda) - \sigma_n(0)}{\lambda} d\lambda.$$

That is

$$\int_0^K \frac{\sigma_n(\lambda) - \sigma_n(0)}{\lambda} d\lambda \leq \log K. \quad (1)$$

Observe that

$$\int_\varepsilon^K \frac{\underline{\sigma}(\lambda) - \bar{\sigma}(0)}{\lambda} d\lambda \leq \int_\varepsilon^K \frac{\liminf_{n \rightarrow \infty} (\sigma_n(\lambda) - \sigma_n(0))}{\lambda} d\lambda.$$

By Fatou's Lemma,

$$\int_\varepsilon^K \frac{\liminf_{n \rightarrow \infty} (\sigma_n(\lambda) - \sigma_n(0))}{\lambda} d\lambda \leq \liminf_{n \rightarrow \infty} \int_\varepsilon^K \frac{\sigma_n(\lambda) - \sigma_n(0)}{\lambda} d\lambda. \quad (2)$$

Since the right hand side of (2) is less than $\log K$, we obtain the following inequality:

$$\int_0^K \frac{\underline{\sigma}(\lambda) - t(0)}{\lambda} d\lambda \leq \log K.$$

Therefore, $\lim_{\lambda \rightarrow 0} \underline{\sigma}(\lambda) = \bar{\sigma}(0)$. That is by Proposition 4.1

$$\bar{\sigma}(0) = t(0). \quad (3)$$

Since one can apply (3) for any subsequence of $\{\sigma_n\}_{n=1}^\infty$ we obtain that

$$\lim_{n \rightarrow \infty} \sigma_n(0) = t(0). \quad \blacksquare$$

5 Spectral convergence

The goal of this section is to prove the main technical proposition of our paper. Note this is based again on the ideas in [5]. Let $P : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a positive, self-adjoint operator and μ_P be its normalized spectral measure that is

$$\mu_P(\lambda) := \frac{\text{the multiplicity of } \lambda \text{ as an eigenvalue}}{n}.$$

Note that if $\|P\| \leq K$ then μ_P is concentrated on the interval $[0, K]$.

Proposition 5.1 *Let $\{Q^n\}_{n=1}^\infty \subset \hat{\Sigma}^d$ be a convergent sequence of oriented simplicial complexes. Then there exists $K > 0$ such that*

- a) *For any $i \geq 1$ and $n \geq 1$, $\|\Delta_{Q^n}^i\| \leq K$.*
- b) *The normalized spectral measures of $\{\Delta_{Q^n}^i\}_{n=1}^\infty$, $\{\mu_n^i\}_{n=1}^\infty$ weakly converge.*
- c) *$c(\mu_n^i) \geq 0$ for any $i, n \geq 1$.*

Proof. To show (a) it is enough to prove the following lemma.

Lemma 5.1 *Let $L, M > 0$ be positive integers. Then if A is a $n \times n$ -matrix of real coefficients (that is a linear operator on $\mathbb{R}^n$) such that*

- each row and column of A contains at most L non-zero elements
- for each entry $A_{i,j}$, $|A_{i,j}| \leq M$

then $\|A\| \leq 2LM$.

Proof. For unit-vectors $f, g \in \mathbb{R}^n$

$$|\langle A(f), g \rangle| = \left| \sum_{1 \leq i, j \leq n} A_{i,j} f(i) g(j) \right| \leq M \sum_{1 \leq i, j \leq n} |f(i) g(j)|.$$

That is

$$|\langle A(f), g \rangle| \leq \sum_{1 \leq i, j \leq n} (f(i)^2 + g(j)^2).$$

Note that the number of occurrences of each $f(i)^2$ or $g(j)^2$ is at most L hence $|\langle A(f), g \rangle| \leq 2LM$. ■

Now let us turn to part (b). The convergence of $\{\mu_n^i\}_{n=1}^\infty$ means that

$$\lim_{n \rightarrow \infty} \int_0^K P(t) d\mu_n^i(t)$$

exists for any real polynomial P . That is one needs to prove that

$$\lim_{n \rightarrow \infty} \frac{\sum_{j=1}^{|Q_i^n|} (\lambda_j^{i,n})^r}{|Q_i^n|}$$

exists where $\{\lambda_j^{i,n}\}$ denotes the spectrum of the i -th Laplacian of Q^n . Hence it is enough to prove that the limit of normalized traces

$$\lim_{n \rightarrow \infty} \frac{\sum_{\sigma \in Q_i^n} (\Delta_{Q_i^n}^i(\sigma, \sigma))^r}{|Q_i^n|} \quad (4)$$

exists. The value of $\Delta_{Q_i^n}^i(\sigma, \sigma)$ depends only on the r -neighborhood of σ , therefore the convergence of the complexes $\{Q^n\}_{n=1}^\infty$ immediately implies the existence of the limit in (4).

Part (c) follows from the simple fact: If Q is a symmetric integer matrix then the product of its non-negative eigenvalues is an integer as well. Indeed, let $\lambda_1, \lambda_2, \dots, \lambda_q$ the list of the non-zero eigenvalues of Q with multiplicities. Let $p(t) = \det(tI - Q)$ be the characteristic polynomial of Q . Then $p(t) = t^s q(t)$, where $q(0) \neq 0$. Obviously, q is an integer polynomial, and $|q(0)| = |\prod_{i=1}^q \lambda_i|$. ■

6 The proof of Theorem 1

We need to prove the following lemma.

Lemma 6.1 *Let $\{K^n\}_{n=1}^\infty \subset \Sigma^d$ be simplicial complexes, then $\lim_{n \rightarrow \infty} \frac{b^i(K_n)}{|V(K^n)|}$ exists for any $i \geq 1$.*

Indeed, by Lemma 6.1 we can immediately see that for any $\varepsilon > 0$ there exists $\delta > 0$ such that if $d_s(K, L) < \delta$ for some $K, L \in \Sigma^d$, then

$$\left| \frac{b^i(K)}{|V(K)|} - \frac{b^i(L)}{|V(L)|} \right| \leq \varepsilon, \quad (5)$$

for any $i \geq 1$. By the definition of the metric d_s there exists some $r \geq 1$ and $\rho > 0$ such that if

$$|p_M(\alpha) - p_N(\alpha)| \leq \rho$$

for any $r' \leq r$ and $\alpha \in Z^{r', d}$ then $d_s(M, N) < \delta$ for any $M, N \in \Sigma^d$.

Now let $L^1, L^2, \dots, L^m$ be a finite set of simplicial complexes such that for any finite simplicial complex K there exists $1 \leq j \leq m$ such that

$$|p_K(\alpha) - p_{L^j}(\alpha)| \leq \frac{\rho}{10} \quad (6)$$

for any $r' \leq r$ and $\alpha \in Z^{r', d}$, where r, ρ are the constants above. The existence of such finite system is clear from compactness.

By the classical Chernoff's inequality there exists $N_\varepsilon > 0$ such that the following holds:

Let $M \in \Sigma^d$ be an arbitrary simplicial complex. Pick N_ε random vertices of M and for any $r' \leq r$ and $\alpha \in Z^{r', d}$ let $Q(M, \alpha)$ be the number of picked vertices x such that $B_{r'}(x) \cong \alpha$. Then

$$\text{Prob} \left\{ \left| \frac{Q(M, \alpha)}{N_\varepsilon} - p_M(\alpha) \right| > \frac{\rho}{10} \text{ for at least one } \alpha \right\} < \varepsilon \quad (7)$$

Thus we have the following testing algorithm. Take the simplicial complex M as an input. Pick N_ε random vertices and calculate $Q(M, \alpha)$ for all $r' \leq r, \alpha \in Z^{r', d}$, where r is the constant above. Check the list $L^1, L^2, \dots, L^m$. By (7), with probability more than $(1 - \varepsilon)$ we find an L^j such that $|Q(M, \alpha) - p_{L^j}(\alpha)| < \frac{\rho}{5}$ for any α . Let $b^i(L_j)$ be our guess. Then by (5) with probability more than $1 - \varepsilon$

$$\left| \frac{b^i(L^j)}{|V(L^j)|} - \frac{b^i(M)}{|V(M)|} \right| < \varepsilon.$$

So, let us prove Lemma 6.1. By Proposition 5.1

$$\lim_{n \rightarrow \infty} \frac{\dim \text{Ker} \Delta_{K^n}^i}{|K_i^n|}$$

exists. Also, by the definition of the convergence of simplicial complexes $\lim_{n \rightarrow \infty} \frac{|K_i^n|}{|V(K_i^n)|}$ exists, hence the lemma, and thus our Theorem follows. $\blacksquare$

References

- [1] I. BENJAMINI AND O. SCHRAMM, *Recurrence of distributional limits of finite planar graphs*. *Electron. J. Probab.* **6** (2001), no. 23, 13 pp. (electronic).
- [2] G. ELEK, *Combinatorial Heat Kernels and Index Theorems*. *Journal of Functional Analysis* **129** (1995) no. 1 64-79.
- [3] G. ELEK, *A Regularity Lemma for Bounded Degree Graphs and Its Applications: Parameter Testing and Infinite Volume Limits* (preprint) URL: <http://arxiv.org/pdf/0711.2800v2>
- [4] O. GOLDBREICH AND D. RON, *Property Testing in Bounded Degree Graphs* *Algorithmica* **32** (2002) 302-343.
- [5] W. LÜCK, Approximating L^2 -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.* **4** (1994), no. 4, 455-481.
- [6] W. LÜCK, L^2 -torsion and 3-manifolds. *Low-dimensional topology (Knoxville, TN, 1992)*, 75-107. Conf. Proc. Lecture Notes Geom. Topology, III, Internat. Press, Cambridge, MA, 1994.